

TB + MYCOLOGY

An extensive selection of Tuberculosis
and Mycology products



Contents

TUBERCULOSIS PRODUCTS	1
Specimen Preparation	1
Stains	2
Direct Specimen Staining Fluorescent Method	3
Reagents & Disks	3
Media for the Cultivation of TB	4
Antimicrobial Susceptibility Testing	6
Control Slides	6
MYCOLOGY PRODUCTS	8
Staining & Slide Preparation	8
Media for the Cultivation of Fungi	9
LAB SUPPLIES	20
Petri Plate Sealing Devices	20
Culture Sampling Device	20
PPE	21

Sabouraud Dextrose Agar
(SabDex), Cat. no. W70R growing
Aspergillus brasiliensis



Specimen Preparation

Digestion and Decontamination Reagents

MucoGest™ 50

MucoGest™ is composed of dithiothreitol, a mucolytic agent recommended for use in the digestion and decontamination procedures of sputum and other clinical specimens for the recovery of mycobacteria and fungi.

0.25 gm (yields 50ml of solution), 20/pk Z130



MucoGest™ for specimen digestion, Cat. no. Z130

TB Prep Kit™

A ready-to-use specimen preparation kit for the processing of sputum specimens for mycobacteria testing. The kit contains five, 50ml TB Base Digestants and five ready-to-use NALC reagents. The components of the kit comply with the CLSI guidelines. Phosphate buffer, for the neutralization step of the procedure, is sold separately.

5 tests/kit Z160

Phosphate Buffer

For the preparation of specimens for AFB culture.

30ml, 2 oz polypropylene bottle, 25/pk X31

40ml, 2 oz polypropylene bottle, 25/pk X43

500ml, polycarbonate bottle, each U10



Phosphate Buffer, Cat. no. U10

Bovine Serum Albumin

Enhances the growth of mycobacteria.

Tube, 3ml, 13x100mm tube,

20/pk Z81

N-Acetyl-L-Cysteine (NALC)

Used in conjunction with TB Base Digestant for specimen preparation for AFB culture.

2.5gm, yields 500ml, 16x100mm tube, 20/pk Z161

250mg, yields 50ml, 16x100mm tube, 20/pk Z60

TB Base Digestant

Used in conjunction with NALC for specimen preparation for AFB culture.

50ml, 2oz polyethylene bottle, 25/pk X45

100ml, 125ml polyethylene bottle, 12/pk U22

500ml, 500ml polypropylene bottle, each U20

Centrifuge Tubes

Screw cap, sterile, polypropylene, printed graduations, 15ml

25/bag, 20 bags per case 6269

Screw cap, sterile, polypropylene, molded graduations, 50ml


25/bag, 20 bags per case 6288



TB Base Digestant, Cat. no. X45

TB PRODUCTS

Kinyoun/Cold Method (Preferred)



Specimen smear (heat or methanol fix).



Carbol Fuchsin 2-3 minutes required (no heat required).



Water rinse.



Use Acid Alcohol Decolorizer until rinsed.



Water rinse.



Counter-stain, Methylene Blue or Brilliant Green.



Water rinse.

Stains

Acid Fast/TB Stains

The Acid-Fast Stain Kit is used as a differential method to detect microorganisms which decolorize with treatment of acid alcohol, and those that do not, such as mycobacteria.

AFB Stain Kit

Carbol Fuchsin (Kinyoun), Acid Alcohol 3%, Methylene Blue Counterstain.

3x8 oz, each

AF900

Acid Alcohol, 3% HCl in Alcohol

Decolorizer for AFB staining, Kinyoun or Ziehl-Neelsen.

250ml, each

AA008

Brilliant Green Counterstain

For Acid-Fast stain procedures and other uses. 1% weight/volume. An aqueous solution of brilliant green.

250ml, each

BG008

AFB Brilliant Green Stain Kit

Contains Carbol Fuchsin (Kinyoun), Acid Alcohol 3%, Brilliant Green Counterstain.

3x8 oz, each

483KB

Carbol Fuchsin for AFB (Kinyoun)

250ml, each

CF008

Methylene Blue Counterstain

For Acid-Fast stain procedures and other uses. A 1% aqueous solution of methylene blue.

16 oz, each

675A16

Methanol (Methyl Alcohol)

32 oz, each

VMT032

1 gallon, each

107B128

473ml bottle, each

QS1016

Acid-Fast Smear Fix

For BACTEC™ blood cultures, for better slide adhesion of specimens.

15ml, each

Z56

Fluorescent Stains

AFB Fluorescent Stain Set (Auramine O)

Auramine O stain kit includes one 250ml bottle of each of the following: auramine O fluorescent stain, decolorizer, and potassium permanganate.
3x250ml, each 418K

Auramine Fluorescent Stain

An aqueous solution of phenol and auramine O Stain.
8 oz, each 418D8

Kit with auramine-rhodamine, fluorescent decolorizer, and potassium-permanganate.
3x250ml, each 630K

Auramine-Rhodamine Stain

An aqueous solution of phenol, glycerine, isopropyl alcohol, rhodamine B, and auramine O.
8 oz, each 789D8

32 oz, each 789D32

Potassium Permanganate (0.55%)

An aqueous solution of potassium permanganate.
8 oz, each PPC18



AFB Fluorescent Stain Set (Auramine O)

Reagents & Disks

Caffeic Acid Disks

A rapid test to detect the ability of an organism to produce the enzyme phenyloxidase; useful for the identification of *Cryptococcus neoformans* and *Cryptococcus gattii*.
25 disks/pk Z118

Nitrate Reagents

Recommended for use in the determination of nitrate utilization by *Mycobacterium* species.

AFB Nitrate Reagent A (Sulfanilimide, 0.2%)
15ml, each Z124

AFB Nitrate Reagent B (Naphthylethylenediamine Dihydrochloride, 0.1%)
15ml, each Z125

Nitrate Reagent C, Zinc Dust

For the completion of the nitrate reduction test. Confirms negatives after adding Nitrate A and Nitrate B reagents.
Each Z73

Hydrochloric Acid, 50% Solution

General purpose laboratory reagent and for use in the *Mycobacterium* nitrate reduction test.
Each Z126



Nitrate Reagent C, Zinc Dust, Cat. no. Z73



Caffeic Acid Disks, for *Cryptococcus*, Cat. no. Z118

Media for the Cultivation of TB



Carbon Utilization Media,
Cat. no. Y125



Lowenstein Jensen Medium,,
Cat. no. C21



HardyFlask™ Lowenstein Jensen
Medium - Gruft, Cat. no. X23

Arylsulfatase Broth

For use in detecting arylsulfatase production in mycobacteria.

0.001M, 2ml, 16x125mm tube, 20/pk K93

Carbon Utilization Media

For the differentiation of *Mycobacterium* spp.

Carbon Utilization Base
Slant, 20x125mm tube, 20/pk Y125

Carbon Utilization with Inositol
Slant, 20x125mm tube, 20/pk Y126

Carbon Utilization with Mannitol
Slant, 20x125mm tube, 20/pk Y127

Carbon Utilization with Citrate
Slant, 20x125mm tube, 20/pk Y128

Lowenstein Jensen Medium

Egg based media for use in the cultivation and isolation of *Mycobacterium* species.

Deep, 20x150mm tube
(for the catalase test), 20/pk C27

Slant, 20x125mm tube, 20/pk C21

HardyFlask™, 20/pk X22

Lowenstein Jensen Medium - Gruft Modification

For the selective isolation of mycobacteria. Contains penicillin, nalidixic acid, and RNA.

Slant, 20x125mm tube, 20/pk C23

HardyFlask™, 20/pk X23

Lowenstein Jensen Medium - Selective

Same formulation as LJ, except for the addition of cycloheximide, lincomycin, and nalidixic acid.

Slant, 20x125mm tube, 20/pk C25

Lowenstein Jensen Medium with Pyruvate

For the cultivation of *Mycobacterium* spp., especially *M. bovis*.

Slant, 20x125mm tube, 20/pk C28

HardyFlask™, 20/pk X19

Lowenstein Jensen with Sodium Chloride

For the differentiation of *Mycobacterium* spp. Contains 5% NaCl.

Slant, 20x125mm tube, 20/pk C29

Lowenstein Jensen with Iron

For the cultivation of *Mycobacterium* spp.

Slant, 20x125mm, 20/pk C37

HardyFlask, 20/pk X21

Middlebrook 7H9

Recommended for use in the cultivation of *Mycobacterium* spp. from sterile fluids.

Middlebrook 7H9 with Tween® 80

5ml, 16x125mm tube, 20/pk C62

Middlebrook 7H9 Broth

10ml, 16x125mm tube, 20/pk C32

Middlebrook 7H10

Agar based media for the isolation and cultivation of *Mycobacterium* species.

Middlebrook 7H10 Agar

26ml, 15x100mm plate, 10/pk W30

Slant, 20x125mm tube, 20/pk C34

HardyFlask™, 20/pk X26

Middlebrook 7H10 with 10% Human Blood

10ml, 20x125mm tube, 20/pk C42

Middlebrook 7H11

Recommended for use in the cultivation and isolation of *Mycobacterium* species.

Middlebrook 7H11 Agar

28ml, 15x100mm plate, 10/pk W35

HardyFlask™, 20/pk X25

Space Saver Plate, 10x100mm, 15/pk SP57

Slant, 20x125mm tube, 20/pk C36

Middlebrook 7H11 Selective

26ml, 15x100mm plate, 10/pk W40

Slant, 20x125mm tube, 20/pk C38

HardyFlask™, 20/pk X28

Middlebrook 7H11/Middlebrook 7H11 Selective

Biplate, 15x100mm, 10/pk J75

Middlebrook OADC Supplement

A supplement for Middlebrook culture media.

100ml, 6 oz bottle, each U98

Nitrate Substrate Broth

For the determination of nitrate reactions of mycobacteria.

2ml, 13x100mm tube, 20/pk R97

Wallenstein Media

Recommended for use in the cultivation and isolation of mycobacteria other than tuberculosis. It is particularly useful in the recovery of *Mycobacteria avium* complex.

Slant, 20x125mm tube, 20/pk C61



Lowenstein Jensen with Iron,
Cat. no. X21



Middlebrook 7H10 Agar,
Cat. no. W30



Middlebrook OADC Supplement,
Cat. no. U98



Antimicrobial Susceptibility Testing

CRITERION™ (Dehydrated Culture Media) Middlebrook 7H10

Recommended for use in qualitative procedures for the isolation and cultivation of mycobacteria species. OADC Supplement, Cat. no. U98, is required for this medium.

Mylar zip-pouch to make 2L	C6280
500 gm wide-mouth bottle	C6281
2kg bucket	C6282
10kg bucket	C6283



CRITERION™ Middlebrook 7H10, 500 gm bottle, Cat. no. C6281

Disk Dispenser

Single metal disk dispenser. Holds one cartridge. Each 260457

For a complete list of antimicrobial susceptibility testing (AST) disks go to HardyDiagnostics.com/Hardy-Disks




Disk Dispenser, Cat. no. 260457

Control Slides

Q-Slides™ AFB Control Slides

Used to test both the stain reagents and user technique. Can be used for Kinyoun, Ziehl-Neelsen, and fluorescent AFB stain methods. Slides are prepared and fixed with a known positive and negative organism. There is also room for test specimens.

5 slides/pk	Z301
-------------	------



Q-Slides™ AFB Control Slide



BlueMount™ - a permanent mounting media with stain included.

Staining & Slide Preparation

BlueMount™

A permanent slide mount for mycology specimens that contains the stain lactophenol cotton blue. Produces a long lasting permanently stained slide. Contains polyvinyl alcohol (PVA) that will dry and set up hard. Creates a perfectly preserved slide.

15ml vial, each Z137

Lactophenol Cotton Blue

For staining fungi on slide wet mounts.

15ml vial, each Z68

Potassium Hydroxide (KOH)

For use in slide preparations of specimens for the purpose of identifying fungal elements. Dissolves epithelial cells that may mask fungal elements.

10%, KOH, 15ml vial, each Z77

20%, KOH, 15ml vial, each Z78

20%, KOH, 40% DMSO, 5ml vial, each Z178



Potassium Hydroxide, 20%, KOH, Cat. no. Z78

India Ink

For use in the identification of *Cryptococcus neoformans* by detection of the polysaccharide capsule. India ink is a negative stain that will show a clear zone where the capsule is.

15ml bottle, each Z64



Media for the Cultivation of Fungi

BiGGY Agar (Nickerson's Agar)

BiGGY Agar (Bismuth Sulfite Glucose Glycine Yeast Agar) is a selective and differential medium used for the isolation and presumptive identification of *Candida* spp.

18ml fill, 15x100mm plate, 10/pk G17

Slant, 13x100mmtube, 20/pk R21

BiGGY Agar, CRITERION™ Dehydrated Culture Media

Mylar® zip-pouch to make 2L C5160

500gm wide-mouth bottle C5161

2kg bucket C5162

10kg bucket C5163



BiGGY Agar, CRITERION™ Dehydrated Culture Media

BHI (Brain Heart Infusion) Agar

Recommended for the cultivation, isolation, and maintenance of yeast and mold.

BHI Agar

26ml, 15x100mm plate, 10/pk W15

Slant, 20x125mm tube, 20/pk L35

Slant, 16x100mm tube, 20/pk L36

HardyFlask™, 20/pk X10

Deep, 20x150mm tube, 20/pk Q16

60ml, 25x100mm plate, 5/pk W163

BHI Agar with Blood

18ml, 15x100mm plate, 10/pk A20

Slant, 20x125mm tube, 20/pk L31

36ml, 25x100mm plate, 5/pk, 50 per box W185BX

HardyFlask™, 20/pk X13

BHI Agar with Blood, Chloramphenicol, and Gentamicin

26ml, 15x100mm plate, 10/pk W65

Slant, 20x125mm tube, 20/pk L114

HardyFlask™, 20/pk X14

BHI Agar with Blood and Gentamicin

HardyFlask™, 20/pk X12

BHI Agar with Chloramphenicol and Cycloheximide

HardyFlask™, 20/pk X11

CGB Agar

For *Cryptococcus gattii*, (canavanine glycine bromothymol blue agar).

20ml, 15x100mm plate, 10/pk G113



BHI Agar with Blood, 15x100mm, Cat. no. A20.



BHI Agar with Blood, Chloramphenicol, and Gentamicin, Cat. no. W65

MYCOLOGY PRODUCTS



Caffeic Acid Agar (Bird Seed Agar), Cat. no. G213

Caffeic Acid Agar (Bird Seed Agar)

Recommended for the selective isolation and differentiation of *Cryptococcus neoformans*. Also known as Bird Seed Agar or Niger Seed Agar.

18ml, 15x100mm plate, 10/pk G213

Caffeic Acid Agar, CRITERION™ Dehydrated Culture Media

For the selective isolation and differentiation of *Cryptococcus neoformans*.

Mylar® zip-pouch to make 2L C7780

500gm wide-mouth bottle C7781

2kg bucket C7782

10kg bucket C7783

Casein Agar

For the differentiation of aerobic actinomycetes based on casein proteolysis. Organisms that can hydrolyse casein, such as *Streptomyces*, and *Pseudomonas*, will produce a clear halo in the surrounding medium.

18ml, 15x100mm plate, 10/pk G122



Cellulose Agar, Modified, CRITERION™ Dehydrated Culture Media, Cat. no. C8730

Cellulose Agar

For the cultivation of fungi from environmental sources.

28ml, 15x100mm plate, 10/pk W83

Cellulose Agar, Modified, CRITERION™ Dehydrated Culture Media

For the cultivation of cellulose-utilizing microorganisms from environmental sources.

Mylar® zip-pouch to make 2L C8730

500gm wide-mouth bottle C8731

2kg bucket C8732

10kg bucket C8733



Czapek Dox Agar, Cat. no. W119

Corn Meal Agar with Tween® 80

Recommended for use in the cultivation of fungi and for the inducement of chlamydo-spore formation by *Candida albicans*.

26ml, 15x100mm plate, 10/pk W10

Czapek Dox Agar

For the cultivation of saprophytic fungi and soil bacteria capable of utilizing inorganic nitrogen. Neutral pH.

28ml fill, 15x100mm plate, 10/pk W119

Czapek Yeast Agar

For the cultivation and maintenance of heat-resistant filamentous fungi from foods or environmental sources.

26ml, 15x100mm plate, 10/pk

W117

Derm-Duet™ II, DTM /RSM (Dermatophyte Test Medium/Rapid Sporulation Media)


For culturing hair, skin and nails for dermatophytes.

DTM will turn red within a few days when a dermatophytic fungus is present. Dermatophytes include *Trichophyton* spp., *Microsporon* spp., and *Epidermophyton* spp. The RSM side will induce the organism to produce spores (macroconidia) which can be viewed microscopically for identification.

RSM media will also turn blue to blue-green in the presence of dermatophytes: making the Derm-Duet™ a perfect combination for the identification of dermatophytic fungi. Individually wrapped.

15x100mm biplate, 10/pk

J175



Derm-Duet™ II, Cat. no. J175

DG-18 Agar (Dichloran-Glycerol)

For the isolation and cultivation of xerophilic molds.

26ml, 15x100mm plate, 10/pk

W85

Dermatophyte Test Medium (DTM)

Dermatophyte Test Medium (DTM) is a selective and differential medium recommended for the cultivation and isolation of pathogenic dermatophytic fungi.

HardyFlask™, 20/pk

X15

Slant, short vial, 20/pk

L27



Dermatophyte Test Medium (DTM), Cat. no. L27

Dermatophyte Milk Agar

Recommended for use in the cultivation and differentiation of certain dermatophytes, particularly *Trichophyton rubrum*, *Trichophyton mentagrophytes*, and *Microsporum persicolor*.

Slant, 16x100mm tube, 20/pk

L75

Dichloran Rose Bengal Chloramphenicol (DRBC) Agar

Recommended for the enumeration of yeast and mold in food and dietary supplements.

18ml, 15x100mm plate, 10/pk

G389

26ml, 15x100mm plate, 10/pk

W89



Dichloran Rose Bengal Chloramphenicol (DRBC) Agar, Cat. no. G389

HardyCHROM™



Candida

18ml, 15x100mm plate, 10/pk

G301

HardyCHROM™ Candida utilizes chromogenic substrates in order to facilitate species identification by color. Due to the unique colors produced, no further testing is needed to identify *C. albicans*, *C. tropicalis*, and *C. krusei*. A trehalose test is needed to confirm *C. glabrata*.


- Helpful in spotting mixed infections.
- Color development occurs within 24 to 48 hours.
- Colors are bright and easy to read out.




C. albicans produces smooth, medium green to dark metallic green colonies.



C. krusei produces rough, spreading, pink colonies.



C. tropicalis colonies appear medium blue to dark metallic blue with a blue halo.



*C. glabrata** produces smooth, pink colonies, often with a darker mauve center.

*Further testing, such as Rapid Trehalose Fermentation Broth (Cat. no. Z205) is needed for confirmation.)

HardyCHROM™

Candida + auris


18ml, 15x100mm plate, 10/pk

G343

HardyCHROM™ *Candida* + *auris* is recommended for the selective isolation and differential identification of *Candida* species. Colonies of *C. auris* will appear white with a characteristic teal to teal-green “bullseye” center and show a unique fluorogenic reaction under UV light at 48-72 hours.


C. auris is unique amongst other *Candida* species because it causes outbreaks and is resistant to nearly all antifungal drugs. This pathogen is also difficult to identify and thus can be misidentified as other species of yeasts. This medium also allows for the differentiation of *C. tropicalis*, *C. albicans* and *C. krusei*, and can aid in the identification of *C. glabrata* when used in conjunction with Rapid Trehalose Broth or GlabrataQuick™. All colonies suspected to be *C. auris* should be subjected to confirmatory identification methods such as MALDI-TOF.

Unique fluorogenic reaction under UV light!



Candida auris at 48 hours under visible ambient light. Note the white perimeter and teal colored center of the isolated colonies.

Candida auris under UV light at 48 hours.




C. tropicalis colonies appear medium blue to dark metallic blue with a blue halo.



C. albicans produces a smooth, medium to green dark metallic green colonies.



C. glabrata produces smooth, pink colonies, often with a darker mauve center.



C. krusei produces rough, spreading, pink colonies.

MYCOLOGY PRODUCTS



Inhibitory Mold Agar (with Chloramphenicol), Cat. no. W58

Inhibitory Mold Agar (with Chloramphenicol)

For the selective isolation of pathogenic fungi.

Contains chloramphenicol for inhibition of bacteria.	
26ml, 15x100mm plate, 10/pk	W25
60ml, 25x100mm plate, 5/pk	W58
HardyFlask™, 20/pk	X20
Slant, 20x125mm tube, 20/pk	L47

Inhibitory Mold Agar (with Gentamicin)

For the selective isolation of pathogenic fungi.

Slant, 20x125mm tube, 20/pk	L49
26ml, 15x100mm plate, 10/pk	W27

Inhibitory Mold Agar, CRITERION™

Dehydrated Culture Media

For the selective isolation of fungi.

Mylar® zip-pouch to make 2L	C5890
500gm wide-mouth bottle	C5891
2kg bucket	C5892
10kg bucket	C5893



m Green Yeast & Fungi Broth, CRITERION™

Dehydrated Culture Media

For the detection of yeast and fungi in beverages using membrane filtration.

Mylar® zip-pouch to make 2L	C7540
500gm wide-mouth bottle	C7541
2kg bucket	C7542
10kg bucket	C7543

Malt Extract Agar with Lecithin and Tween, Cat. no. P93

Malt Extract Agar with Lecithin and Tween

For the cultivation, isolation, and maintenance of yeast and mold.

26ml, 15x100mm plate, 10/pk	W28
-----------------------------	-----

Malt Extract Agar with Lecithin and Tween

For the cultivation of yeasts and molds from environmental surfaces.

15x65mm contact plate, 10/pk	P93
------------------------------	-----

Malt Extract Agar with 0.01% Chloramphenicol

For the selective isolation of fungi.

26ml, 15x100mm plate, 10/pk	W80
Slant, 16x100mm tube, 20/pk	L128



Aspergillus on Malt Extract Agar, Cat. no. W80

Nocardia Hydrolysis Kit for Actinomycetes

Nocardia media are designed to aid in the differentiation of aerobic Actinomycetes, *Nocardia*, *Actinomadura*, and *Streptomyces* species by determining the presence of hydrolysis enzymes. Box contains 5 each of Casein, Xanthine, Hypoxanthine, and Tyrosine Agars.

Deep, 20x125mm tubes, 20/pk Z604

Osmophilic Yeast Agar Plate

Recommended for use in the enumeration of osmophilic yeast in food.

15x100mm plate, 10/pk G199



Osmophilic Yeast Agar Plate, Cat. no. G199

Potato Dextrose Agar

Recommended for use in the cultivation of yeast and mold and for the inducement of sporulation.

26ml, 15x100mm plate, 10/pk W60

68ml, 15x150mm plate, 10/pk H51

18ml, 20x125mm tube, 20/pk Q29

200ml, 250ml glass bottle, 12/pk U294

Slant, 10ml, 20x125mm tube, 20/pk L90

Potato Dextrose Agar with Chlortetracycline

For use in the cultivation of yeast and mold and for the inducement of sporulation.

26ml, 15x100mm plate, 10/pk W93

Potato Dextrose Agar with TA (Tartaric Acid)

For determining yeast and mold counts from butter, margarine, and related products.

13ml, 15x60mm plate, 10/pk G94

28ml, 15x100mm plate, 10/pk W96

Potato Dextrose Agar, CRITERION™ Dehydrated Culture Media

For the cultivation and identification of fungi.

Mylar® zip-pouch to make 2L C6620

500g wide-mouth bottle C6621

2kg bucket C6622

10kg bucket C6623

Potato Flake Agar

Used for the cultivation, isolation, and induction of sporulation of fungi from clinical specimens.

26ml, 15x100mm plate, 10/pk W59

HardyFlask™, 20/pk X32



Potato Dextrose Agar, Cat. no. U294



Potato Dextrose Agar with TA (Tartaric Acid), Cat. no. G94

MYCOLOGY PRODUCTS



Rose Bengal Chloramphenicol Agar, Cat. no. W87

Potato Flake Selective Agar

Potato Flake Selective Agar additionally contains Ciprofloxacin and Vancomycin.

26ml, 15x100mm plate, 10/pk	W159
HardyFlask™, 20/pk	X59

Rose Bengal Agar with Chloramphenicol

For the enumeration of yeasts and molds in food and the environment.

24ml, 15x100mm plate, 10/pk	W87
18ml, 20x150mm tube, 10/pk	Q81
15x65mm contact plate, 10/pk	P42

Rose Bengal Agar Base, CRITERION™ Dehydrated Culture Media

For the enumeration of yeasts and molds in food and the environment.

500gm wide-mouth bottle	C8101
2kg bucket	C8102



Sabdex (Sabouraud Dextrose) Agar, Cat. no. U270

Sabouraud Dextrose Agar

Recommended for the isolation, cultivation, and maintenance of non-pathogenic and pathogenic fungi and yeast.

15x65mm contact plate, 10/pk	P36
26ml, 15x100mm plate, 10/pk	W70
26ml, 15x100mm Red Tinted plate, 10/pk	W70R
70ml, 15x150mm plate, 10/pk	H39
60ml, Deep Fill, 25x100mm plate, 5/pk	W250
HardyFlask™, 50ml plastic, 20/pk	X40

Tube

Deep, 20x125mm, 20/pk	Q31
Deep, 20x150mm, USP, 10/bx	Q83
Slant, 20x125mm, 20/pk	L40

Bottle

200ml, 500ml polycarbonate, 10/pk	U270
200ml, 16 oz glass boston round, 12/pk	U227
400ml, 16 oz glass, 12/pk	U352
500ml, 500ml polycarbonate, 10/pk	U353

Sabouraud Dextrose Agar (with Chloramphenicol and Tetracycline)

For the selective isolation of fungi.

26ml, 15x100mm plate, 10/pk	W74
-----------------------------	-----



Sabouraud Dextrose Agar (with Chloramphenicol and Tetracycline), Cat. no. W74

Sabouraud Dextrose Agar with Lecithin and Tween® 80

For the isolation, cultivation, and maintenance of non-pathogenic and pathogenic species of fungi and yeasts.

15ml, 15x65mm contact plate, 10/pk	P46
26ml, 15x100mm plate, 10/pk	W71
400ml fill, 16 oz glass bottle, 12/pk	U431
400ml fill, 500ml polycarbonate bottle, 10/pk	U175
Deep, 20x125mm tube, 20/pk	Q75

Sabouraud Dextrose Agar, USP, Blue Tint

Tinted blue for easy differentiation from TSA.

26ml fill, 15x100mm plate, 10/pk	W82
----------------------------------	-----

Sabouraud Dextrose Agar (with Chloramphenicol)

For the selective isolation of fungi and yeasts from clinical and non-clinical specimens.

26ml, 15x100mm plate, 10/pk	W72
20x125mm tube, 20/pk	Q34
HardyFlask™, 20/pk	X41

Sabouraud Dextrose Agar, Emmon's, CRITERION™ Dehydrated Culture Media

For the isolation, cultivation, and maintenance of non-pathogenic and pathogenic species of fungi and yeasts.

Mylar® zip-pouch to make 2L	C6770
500gm wide-mouth bottle	C6771
2kg bucket	C6772
10kg bucket	C6773

Sabouraud Dextrose Agar with Chloramphenicol and Gentamicin

For the selective isolation of fungi.

12ml, 15x60mm plate, 10/pk	G159
26ml, 15x100mm plate, 10/pk	W73

Sabouraud Dextrose (with Chloramphenicol and Gentamicin) / DTM

For the selective isolation and differentiation of pathogenic fungi and dermatophytic fungi.

15x100mm biplate, 10/pk	J107
-------------------------	------



Sabouraud Dextrose Agar with Lecithin and Tween® 80, Cat. no. U431



Sabouraud Dextrose Agar, Blue Tint, Cat. no. W82



Sabouraud Dextrose (with Chloramphenicol and Gentamicin) / DTM, Cat. no. J107

MYCOLOGY PRODUCTS



Sabouraud Dextrose Agar, Emmon's, Cat. no. X57

Sabouraud Dextrose Agar, Emmon's

For the isolation, cultivation, and maintenance of non-pathogenic and pathogenic species of fungi and yeasts.

26ml, 15x100mm plate, 10/pk	W20
HardyFlask™, 20/pk	X57

Sabouraud Dextrose Agar, CRITERION™ Dehydrated Culture Media

For the isolation, cultivation, and maintenance of non-pathogenic and pathogenic species of fungi.

Mylar® zip-pouch to make 2L	C6810
500gm wide-mouth bottle	C6811
2kg bucket	C6812
10kg bucket	C6813



Sabouraud Dextrose Agar (with Chloramphenicol), CRITERION™, Cat. no. C6783

Sabouraud Dextrose Agar (with Chloramphenicol), CRITERION™ Dehydrated Culture Media

A selective agar medium for the isolation, identification and cultivation of fungi.

Mylar® zip-pouch to make 2L	C6780
500gm wide-mouth bottle	C6781
2kg bucket	C6782
10kg bucket	C6783

Sabouraud Dextrose Agar with Lecithin and Tween® 80, CRITERION™ Dehydrated Culture Media

For the cultivation of fungi from environmental surfaces.

Mylar® zip-pouch to make 2L	C6800
500gm wide-mouth bottle	C6801
2kg bucket	C6802
10kg bucket	C6803



Sabouraud Dextrose Broth, CRITERION™ Dehydrated Culture Media, Cat. no. C7562

Sabouraud Dextrose Broth, CRITERION™ Dehydrated Culture Media

For the isolation, cultivation, and maintenance of non-pathogenic and pathogenic species of fungi and yeasts.

Mylar® zip-pouch to make 2L	C7560
500gm wide-mouth bottle	C7561
2kg bucket	C7562
10kg bucket	C7563

Sabouraud Dextrose (SabDex) Broth

For the isolation, cultivation, and maintenance of non-pathogenic and pathogenic species of fungi and yeasts.

10ml fill, 16x125mm tube, 20/pk K54

SabHI Agar

A combination of SabDex and Heart Infusion Agar. Recommended for use in the cultivation of fungi.

26ml, 15x100mm plate, 10/pk W75

HardyFlask™, 20/pk X75

SabHI Agar with Blood, Chloramphenicol, and Cycloheximide

For the selective isolation of pathogenic fungi.

HardyFlask™, 20/pk X73

SabHI Agar Base, CRITERION™ Dehydrated Culture Media

For the isolation and cultivation of fungi.

Mylar® zip-pouch to make 2L C6790

500gm wide-mouth bottle C6791

2kg bucket C6792

10kg bucket C6793

Trehalose Broth, Rapid

A rapid screening method to aid in the detection of *Candida glabrata*.

Tube, 0.25ml, 13x100mm tube, 20/pk Z205

Trichophyton Agar

Recommended for the cultivation and differentiation of *Trichophyton* spp.

Tube

Trichophyton #1, 10.5ml, 20x125mm, 20/pk C51

Trichophyton #2, 10.5ml, 20x125mm, 20/pk C52

Trichophyton #3, 10.5ml, 20x125mm, 20/pk C53

Trichophyton #4, 10.5ml, 20x125mm, 20/pk C54

Trichophyton #5, 10.5ml, 20x125mm, 20/pk C55

V9 Agar (V8 juice and Potato Flakes)

Recommended for the rapid growth and sporulation of fungi.

22ml fill, 15x100mm plate, 10/pk G98



SabHI Agar with Blood, Chloramphenicol, and Cycloheximide, Cat. no. X73



Trehalose Broth, Rapid, Cat. no. Z205



V9 Agar (V8 juice and Potato Flakes), Cat. no. G98



MycoSeal™ Clear, Shrink Seals

Petri Plate Sealing Devices

MycoSeal™ Plate Sealing Bands

MycoSeal™ is a cellulose shrink seal designed to keep Petri plate lids tightly sealed. Sealed plates retain their moisture and stay fresh longer. Ideal for mycology and TB cultures that require long incubation times. MycoSeal™ prevents dehydration and cracking of the media and yet still allows both oxygen and carbon dioxide gas exchange. For 100mm plates only.

60 per/pk

SS9225



PetriSeal™ Stretch Tape

PetriSeal™ Stretch Tape

Sealing tape specifically designed for the special requirements of Petri plate, microtiter plates and culture plates and their corresponding lids. The unique elastomeric characteristics allow the tape to stretch and conform around the irregular perimeters of these vessels to form a tight seal. The seal will block water vapor loss, reduce the rate of gas exchange, and prevent the entry of contaminants. Waterproof and chemically resistant to most organic solvents and caustic agents. The seal remains flexible down to -30°C and withstands continuous exposure from -40°C to + 77°C (with intermittent exposure up to 100°C). Heavy 5mil (0.005in.) thick tape is durable and resists abrasion and puncture.

White, 0.5 inches x 108 feet

PSEAL108W

Culture Sampling Device

MycoMount™

MycoMount™ is a device used to facilitate the preparation of fungal wet mounts for microscopic examination. A sticky adhesive pad on a long handle allows easy access to tubes and bottles and eliminates the need for “tease mounts” and “scotch tape preps.” After collection of fungal elements the pad can be mounted directly on a microscope slide for convenient viewing.

40 strips/pk

MM40



MycoMount™

PPE

ComfortPRO™ Lab Coat

These affordable, disposable lab coats are made of tri-laminated, spunbound material (SMS) which is designed to be fully breathable and splash resistant.

- Knit cuffs and collars for added comfort
- Three pockets are featured, one on the chest and two at the hip, with snap front buttons

Medium, 70/pk NEGU6073M

Large, 70/pk NEGU6073L

XLarge, 70/pk NEGU6073XL



ComfortPRO™ Lab Coat

Lab Apron

Personal protective apron made of polyethylene, 2-1/4 mil. Perfect for processing specimens where splashing may occur. Plastic, fluid proof.

43"x59", 20/pk 8571

Face Shield, Full Face, Disposable

Face shields are personal protective equipment (PPE) that are used for protection of the facial areas (eyes, nose, mouth) from sprays, aerosols, and splashes.

- Well suited for protection against COVID-19 and other pathogens
- Ideal for healthcare and laboratory workers
- Made from PETG
- Provides clarity of view
- Visors are anti-fogging
- One size fits most

10/pk MD30331



Face Shield, Full Face, Disposable

Splash Shield

A lightweight and comfortable face shield that is optically clear, has anti-fog properties, fits over glasses, a foam headband, and is disposable/reusable. Designed to protect the user's face from splattered blood and other bodily fluids, when used properly.

Splash Shield for face, 24/pk 4511



Splash Shield

Precision® Nitrile, Exam Gloves

Precision® Violet Blue Nitrile Powder-Free Exam Gloves are ultra soft, and super stretchy to reduce hand fatigue and maximize comfort. 100% synthetic and latex free with a textured surface to provide strong grip in both dry and wet applications. Thinner for maximum dexterity but very strong to provide superior barrier protection.

Small, 1,000/bx PCS772

Medium, 1,000/bx PCS775

Large, 1,000/bx PCS776

Extra Large, 1,000/bx PCS778



Exam Gloves, Cat. no. PCS775



HARDY DIAGNOSTICS

A Culture of Service™

40
MORE THAN



Years of
Bringing You
A Culture of Service



100%
Employee Owned

ESTABLISHED

1980

MADE IN
USA



Hardy Diagnostics
donates 1% of net
profits to charity.



FM 572526

Hardy Diagnostics has a Quality Management System that is certified to ISO 13485 and is a FDA licensed medical device manufacturer.

Headquarters

1430 West McCoy Lane
Santa Maria, CA 93455
800.266.2222
Sales@HardyDiagnostics.com
HardyDiagnostics.com

Distribution Centers

Santa Maria, California
Olympia, Washington
Salt Lake City, Utah
Phoenix, Arizona
Dallas, Texas
Des Moines, Iowa
Springboro, Ohio
Lake City, Florida
Albany, New York
Raleigh, North Carolina



Contract Holder

Contract # 47QSWA22D0077